

NEW AND IMPROVED BLUE GEL For Use by Licensed Professionals Only

This product contains 5% lidocaine and 1% tetracaine, together with a vascular constrictor which reduces swelling, bruising, and bleeding. This is the anesthetic of choice for fast onset and reduction of swelling in areas such as lips or inner arms, where bruising and bleeding are a problem.

FOR EXTERNAL USE ONLY: Temporarily relieves pain and swelling associated with tattooing, permanent makeup, piercing or other pain sensitive procedures. Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool, dark place, *out of reach of children.*

Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline, and seek medical attention. **DO NOT SWALLOW.** If you should accidentally swallow this product, seek medical attention immediately. Do not use if pregnant or nursing.

DURING PROCEDURE ANESTHETIC

2-5 minutes

SUGGESTED APPLICATION

Use this product during painful procedures to continue the anesthetic effects of your pre-procedure topical anesthetics, or to deaden broken skin. *It will have no effect if applied to intact epidermis.*

Always wear gloves while applying. Apply to the procedure area with your gloved hand or an appropriate applicator. Cover with plastic and wait 2-5 minutes or until anesthetic effect occurs.

Rinse off thoroughly before continuing your procedure so that the product does not penetrate the skin. **DO NOT REAPPLY THIS PRODUCT TO THE SAME AREA** because it can cause the skin to become so anesthetized that you could harm the tissue. If you need more anesthetic effect before you have finished your procedure, use **Top Shelf Super T** to continue the anesthetic effect.

New and Improved Blue Gel or **Five-Star Vasocaine** will keep the skin anesthetized and constricted for up to two hours.

NEW AND IMPROVED BLUE GEL For Use by Licensed Professionals Only

This product contains 5% lidocaine and 1% tetracaine, together with a vascular constrictor which reduces swelling, bruising, and bleeding. This is the anesthetic of choice for fast onset and reduction of swelling in areas such as lips or inner arms, where bruising and bleeding are a problem.

FOR EXTERNAL USE ONLY: Temporarily relieves pain and swelling associated with tattooing, permanent makeup, piercing or other pain sensitive procedures. Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool, dark place, *out of reach of children.*

Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline, and seek medical attention. **DO NOT SWALLOW.** If you should accidentally swallow this product, seek medical attention immediately. Do not use if pregnant or nursing.

DURING PROCEDURE ANESTHETIC

2-5 minutes

SUGGESTED APPLICATION

Use this product during painful procedures to continue the anesthetic effects of your pre-procedure topical anesthetics, or to deaden broken skin. *It will have no effect if applied to intact epidermis.*

Always wear gloves while applying. Apply to the procedure area with your gloved hand or an appropriate applicator. Cover with plastic and wait 2-5 minutes or until anesthetic effect occurs.

Rinse off thoroughly before continuing your procedure so that the product does not penetrate the skin. **DO NOT REAPPLY THIS PRODUCT TO THE SAME AREA** because it can cause the skin to become so anesthetized that you could harm the tissue. If you need more anesthetic effect before you have finished your procedure, use **Top Shelf Super T** to continue the anesthetic effect.

New and Improved Blue Gel or **Five-Star Vasocaine** will keep the skin anesthetized and constricted for up to two hours.